

Phospho-ADAM9 (Thr761) Antibody

| | | |
|-------------------|----------------|-----------------------|
| Cat.#: AF7059 | Concn.: 1mg/ml | Mol.Wt.: 91kDa,105kDa |
| Size: 100ul,200ul | Source: Rabbit | Clonality: Polyclonal |

Application: WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000

Reactivity: Human,Mouse,Rat


Purification: The antibody is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.

Specificity: Phospho-ADAM9 (Thr761) Antibody detects endogenous levels of ADAM9 only when phosphorylated at Thr761.

Immunogen: A synthesized peptide derived from human ADAM9 around the phosphorylation site of Thr761.


Uniprot: Q13443

Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.




Western blot analysis of extracts from RAW264.7 cells(H2O2 treatment), using Phospho-ADAM9 (Thr761) Antibody. The lane on the left was treated with blocking peptide.


Observed bands: 105kDa




Western blot analysis of extracts from Mouse kidney tissue, using Phospho-ADAM9(Thr761) Antibody. Lane1 was treated with phospho-blocking peptide, Lane2 was treated with non-phospho-blocking peptide.



Western blot analysis of extracts from Mouse kidney tissue sample, using Phospho-ADAM9(Thr761) Antibody(AF7059).



AF7059 at 1/100 staining mouse kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.



AF7059 at 1/100 staining human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary antibody.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.