

ADIPOR1 Ab

Cat.#: DF6430
Size: 100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 43kDa
Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200

Reactivity: Human,Mouse,Rat

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Specificity: ADIPOR1 Ab detects endogenous levels of total ADIPOR1.

Immunogen: A synthesized peptide derived from human ADIPOR1.

Uniprot: Q96A54

Description: Adiponectin is a circulating hormone secreted by adipocytes that improves the metabolism of glucose and lipids, and is expressed at low levels in those with obesity and diabetes. Adiponectin receptors AdipoR1 and AdipoR2, also designated progestin and adipoQ receptor family members I and II, respectively, regulate fatty acid oxidation and the uptake of glucose by adiponectin. Each receptor activates a unique set of signaling molecules including AMPK, p38 MAPK and PPAR . AdipoR1 has a high-affinity for globular adiponectin and low-affinity for full-length adiponectin, while AdipoR2 has an intermediate affinity for both forms. AdipoR1 and AdipoR2 are mainly expressed in liver and muscle. Adiponectin, AdipoR1 and AdipoR2 are all associated with body composition, Insulin sensitivity, and metabolic parameters. Physical training increases circulating adiponectin and mRNA expression of AdipoR1 and AdipoR2 in muscle, which may mediate the improvement of Insulin resistance and the metabolic syndrome in response to exercise.

Subcellular Location: Membrane. Localized to the cell membrane and intracellular organelles.


Tissue Specificity: Widely expressed (PubMed:16044242). Highly expressed in heart and skeletal muscle (PubMed:12802337). Expressed at intermediate level in brain, spleen, kidney, liver, placenta, lung and peripheral blood leukocytes (PubMed:12802337). Weakly expressed in colon, thymus and small intestine (PubMed:12802337).

Similarity: The N-terminus is cytoplasmic and the C-terminus is extracellular, contrary to what is observed for G-protein

coupled receptors. Unlike G-protein coupled receptors, transmembrane helices are not kinked or tilted relative to the plane of the membrane. Belongs to the ADIPOR family.

Storage Condition and Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.



Western blot analysis of extracts of Y79, using ADIPOR1 Ab. The lane on the left is treated with the antigen-specific peptide.



DF6430 at 1/100 staining Human lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.