

## CDT1 Ab

Images(3)

Cat.#: DF8171  
Size:

Concn.: ~1mg/ml  
Source: Rabbit

Mol.Wt.: 63 kDa  
Clonality: Polyclonal

Application: IF/ICC 1:100-1:500, WB 1:1000-3000


\*The optimal dilutions should be determined by the end user.  
Reactivity: Human,Mouse,Rat

Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.


Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human CDT1, corresponding to a region within C-terminal amino acids.

Uniprot: Q9H211



Western blot analysis of extracts from Rat brain, using CDT1 Ab. The lane on the left was treated with blocking peptide.



DF8171 staining A549 cells(UV treatment) by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(#DF8171) and mouse anti-beta tubulin Ab(#T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.