

FAM111B Ab

Images(4)

Cat.#: DF8839
Size:

Concn.: ~1mg/ml
Source: Rabbit

Mol.Wt.: 84 kDa
Clonality: Polyclonal

Application:

IF/ICC 1:100-1:500, IHC 1:50-1:200, WB 1:1000-3000

*The optimal dilutions should be determined by the end user.
Reactivity:

Human,Mouse

Storage:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification:


The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen:


A synthesized peptide derived from human F111B, corresponding to a region within N-terminal amino acids.

Uniprot:


Q6SJ93



Western blot analysis of extracts from various samples, using FAM111B Ab.
Lane 1: Mcf7 cells(serum starvation treatment), blocked with antigen-specific peptides.
Lane 2: Mcf7 cells(serum starvation treatment).
Lane 3: K562 cells(uv treatment).



DF8839 at 1/100 staining rat heart tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-rabbit Ab was used as the secondary Ab.



DF8839 staining A549 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(#DF8839) and mouse anti-beta tubulin Ab(#T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG Ab(Green) were used as the secondary Ab.
The nuclear counter stain is DAPI (blue).



Affinity Biosciences
website:www.affbiotech.com
order:order@affbiotech.com

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.