Phospho-CD31 (Tyr713) Ab

Images(3)

Cat.#: AF3191 Mol.Wt.: 82/150kDa Concn.: ~1mg/ml Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Human, Mouse Reactivity:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

A synthesized peptide derived from human CD31 around the Immunogen:

phosphorylation site of Tyr713.

P16284 Uniprot:

Description: PECAM-1 is a cell adhesion molecule expressed on platelets and at

endothelial cell intercellular junctions.




Western blot analysis of extracts from various samples, using Phospho-PECAM-1 (Tyr713) Ab.

Lane 1: UV treated MDA-MB-231 cells, blocked with antigen-specific

peptides,

Lane 2: UV treated MDA-MB-231 cells,

Lane 3: H2O2 treated EC304 cells.



AF3191 staining H2O2 treated Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF3191) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI(blue).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.



For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.