

Phospho-BLNK (Tyr96) Ab

[Images\(2\)](#)

Cat.#: AF3467
 Size:

Concn.: ~1mg/ml
 Source: Rabbit

Mol.Wt.: 50kDa
 Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
 *The optimal dilutions should be determined by the end user.

Reactivity: Human,Mouse


Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho-peptide and non-phospho-peptide affinity columns.


Immunogen: A synthesized peptide derived from human BLNK around the phosphorylation site of Tyr96.

Uniprot: Q8WV28

Description: BLNK an adaptor protein that bridges the B-cell receptor-associated kinases (BCR) with a multitude of signaling pathways, regulating biologic outcomes of B-cell function and development. Plays an important role in BCR-mediated PLCG1 activation and Ca(2+) mobilization.



Western blot analysis of Phospho-BLNK (Tyr96) Ab expression in COLO205 cells lysates. The lane on the right was treated with the antigen-specific peptide.



AF3467 at 1/100 staining human lymph node tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking,



Affinity Biosciences
website:www.affbiotech.com
order:order@affbiotech.com

overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.