

## **IGF1 Receptor Ab**

References(4) Images(4)

Cat.#: AF6123 Concn.: ~1mg/ml Mol.Wt.: 90,155kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink<sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human IGF1 Receptor, corresponding to


a region within C-terminal amino acids.

Uniprot: P08069

Description: This receptor binds insulin-like growth factor with a high affinity. It has

tyrosine kinase activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates

alpha and beta subunits.



Western blot analysis of extracts from K562, using IGF1R Ab. The lane on

the left was treated with blocking peptide.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.