

## **Affinity Biosciences** website:www.affbiotech.com order:order@affbiotech.com

## CD44 Ab

References(2) Images(4)

Cat.#: AF6186 Mol.Wt.: 90kDa Concn.: ~1mg/ml Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

A synthesized peptide derived from human CD44, corresponding to a region Immunogen:

within C-terminal amino acids.

P16070 Uniprot:

Description: The protein encoded by this gene is a cell-surface glycoprotein involved in

> cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs).

25 — 20

150

the left was treated with blocking peptide.

AF6186 staining HepG2 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF6186) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

Western blot analysis of extracts from 293 cells, using CD44 Ab. The lane on

The nuclear counter stain is DAPI(blue).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.



For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.