Phospho-ATM (Ser1981) Ab

References(2) Images(5)

Cat.#: AF4120 Concn.: ~1mg/ml Mol.Wt.: 350kd Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:1000, IHC 1:50-1:200

*The optimal dilutions should be determined by the end user.

Reactivity: Human

Storage: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and

Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store

at -20 °C. Stable for 12 months from date of receipt.

Purification: The Ab is from purified rabbit serum by affinity purification via sequential

chromatography on phospho-peptide and non-phospho-peptide affinity

columns.

Immunogen: Peptide sequence around phosphorylation site of serine 1981 (E-G-S(p)-Q-


S) derived from Human ATM.

Uniprot: Q13315

Description: ATM encoded by this gene belongs to the PI3/PI4-kinase family. This

protein is an important cell cycle checkpoint kinase that phosphorylates; thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability. Mutations in this gene are associated with ataxia telangiectasia, an autosomal recessive disorder. Two transcript variants encoding different

isoforms have been found for this gene.



AF4120 at 1/100 staining human liver carcinoma sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.



For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.