


## ATF5 Ab

[References\(1\)](#) [Images\(2\)](#)

|               |                                                                                                                                                               |                       |
|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| Cat.#: DF3480 | Concn.: ~1mg/ml                                                                                                                                               | Mol.Wt.: 31 KD        |
| Size:         | Source: Rabbit                                                                                                                                                | Clonality: Polyclonal |
| Application:  | WB 1:500-1:1000, IF/ICC 1:100-1:500<br>*The optimal dilutions should be determined by the end user.                                                           |                       |
| Reactivity:   | Human, Mouse, Rat                                                                                                                                             |                       |
| Storage:      | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt. |                       |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).                                     |                       |
| Immunogen:    | A synthesized peptide derived from human ATF5, corresponding to a region within N-terminal amino acids.                                                       |                       |
| Uniprot:      | Q9Y2D1                                                                                                                                                        |                       |



DF3480 staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF3480) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.