

CD20 Ab

Cat.#: BF0233	Concn.: ~1mg/ml	Mol.Wt.: 33kDa.
Size:	Source: Mouse	Clonality: Monoclonal

Application: ELISA 1:10000, IHC 1:200-1:1000, IF/ICC 1:200-1:1000
*The optimal dilutions should be determined by the end user.

Reactivity: Human

Storage: Mouse IgG1 in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification: Affinity-chromatography.

Immunogen: Purified recombinant fragment of human CD20 expressed in E. Coli.

Uniprot: P11836

Description: CD20 (MS4A1, membrane-spanning 4-domains, subfamily A, member 1) is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. The CD20 antigen is present on human pre B lymphocytes and on B lymphocytes at all stages of maturation, except on plasma cells. Low level expression of the CD20 antigen has been detected on normal T lymphocytes. The CD20 molecule is involved in regulation of B cell differentiation, presumably via its reported function as a Ca⁺⁺ channel subunit. And it is known to accelerate the G0 to G1 progression induced by IGF-1.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.