Myeloperoxidase Ab

Images(1)

Cat.#: BF0417 Concn.: ~1mg/ml Mol.Wt.: 84kDa Size: Source: Mouse Clonality: Monoclonal

Application: ELISA 1:10000, WB 1:500-1:2000

*The optimal dilutions should be determined by the end user.

Reactivity: Human

Storage: Mouse IgG1 in phosphate buffered saline (without Mg2+ and Ca2+), pH

7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C.

Stable for 12 months from date of receipt.

Purification: Affinity-chromatography.

Immunogen: Purified recombinant fragment of human Myeloperoxidase expressed in E.

Coli.

Uniprot: P05164

Description: Myeloperoxidase (MPO) is a heme protein synthesized during myeloid

differentiation that constitutes the major component of neutrophil

azurophilic granules. Produced as a single chain precursor, myeloperoxidase is subsequently cleaved into a light and heavy chain. The mature myeloperoxidase is a tetramer composed of 2 light chains and 2 heavy

chains. This enzyme produces hypohalous acids central to the microbicidal

activity of netrophils.




Figure 1: Western blot analysis using MPO mouse mAb against truncated Trx-MPO recombinant protein (1),truncated MBP-MPO (aa1-193) recombinant protein (2) and truncated MPO(aa165-745)-hIgGFc transfected CHO-K1 cell

lysate(3).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.