

NTF3 Ab

[References\(3\)](#) [Images\(5\)](#)

Cat.#: DF6105	Concn.: ~1mg/ml	Mol.Wt.: 29kDa
Size:	Source: Rabbit	Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human, Mouse, Rat	
Storage:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human NTF3, corresponding to a region within the internal amino acids.	
Uniprot:	P20783	
Description:	NT-3 is a member of the structurally related neurotrophin family of proteins, which includes ?-NGF, BDNF and NT-4. NT-3 is expressed in a number of cell types including neuronal cells, eosinophils, and melanocytes (1-3). NT-3 is required for the development of peripheral sensory neurons. NT-3 is secreted from cells as a precursor protein, which is proteolytically cleaved into the mature form. NT-3 signaling is mediated through two distinct receptors, the neurotrophin receptor p75NTR and the Trk tyrosine kinase receptor TrkC. While all neurotrophins bind to the p75NTR receptor, NT-3 preferentially binds to TrkC and, with lesser affinity, TrkB.	

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.