

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

PARK7 Ab

Images(2)

Cat.#: DF6169 Concn.: ~1mg/ml Mol.Wt.: 20kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human PARK7, corresponding to a

region within C-terminal amino acids.

Uniprot: Q99497

Description: Parkinson's disease (PD) is characterized by the presence of Lewy bodies

(intracellular inclusions) and by the loss of dopaminergic neurons. Research studies have shown that mutations in ?-synuclein, Parkin, and DJ-1 are linked to PD . ?-synuclein is a major component of the aggregates found in Lewy bodies. Parkin is involved in protein degradation through the ubiquitin-proteasome pathway, and investigators have shown that mutations in Parkin cause early onset of PD . Loss-of-function mutations in DJ-1 cause early onset of PD, but DJ-1 is associated with multiple functions: it cooperates with Ras to increase cell transformation, it positively regulates transcription of the androgen receptor, and it may function as an indicator of oxidative

stress (2-5).



Western blot analysis of extracts from HepG2, using PARK7 Ab.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



DF6169 at 1/100 staining Rat heart tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4° C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.