

website:www.affbiotech.com order:order@affbiotech.com

IL24 Ab

Images(2)

Cat.#: DF6672 Mol.Wt.: 24kDa Concn.: ~1mg/ml Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

A synthesized peptide derived from human IL24, corresponding to a region Immunogen:

within the internal amino acids.

O13007 Uniprot:


Description: This gene encodes a member of the IL10 family of cytokines. It was

> identified as a gene induced during terminal differentiation in melanoma cells. The protein encoded by this gene can induce apoptosis selectively in various cancer cells. Overexpression of this gene leads to elevated expression of several GADD family genes, which correlates with the induction of apoptosis. The phosphorylation of mitogen-activated protein kinase 14 (MAPK7/P38), and heat shock 27kDa protein 1 (HSPB2/HSP27) are found to be induced by this gene in melanoma cells, but not in normal immortal melanocytes. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Western blot analysis of A431 cell lysates(including supernates), using IL24 Ab. The lane on the left was treated with the antigen-specific peptide.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



DF6672 at 1/100 staining Human kidney cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4° C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.