

TNFRSF14 Ab

Images(2)

Cat.#: DF6724

Size:

Concn.: ~1mg/ml

Source: Rabbit

Mol.Wt.: 30kDa

Clonality: Polyclonal

Application:

WB 1:500-1:2000, IHC 1:50-1:200

*The optimal dilutions should be determined by the end user.

Reactivity:

Human,Rat

Storage:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.

Purification:

The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen:


A synthesized peptide derived from human TNFRSF14, corresponding to a region within C-terminal amino acids.

Uniprot:


Q92956

Description:

The mammalian tumor necrosis factor receptor (TNFR) family consists of 10 cell-surface proteins that regulate development and homeostasis of the immune system. Member 14 of TNFR (TNFRSF14, also named as TR2, ATAR, HVEA, HVEM, LIGHTR) was also identified as a mediator of herpesvirus entry into mammalian cells. The cytoplasmic region of TNFRSF14 bound to several members of the TNFR-associated factor (TRAF) family, namely TRAF1, TRAF2, TRAF3, and TRAF5. Transient transfection into human 293 cells caused marked activation of nuclear factor- B (NF- B), a transcriptional regulator of multiple immunomodulatory and inflammatory genes.



Western blot analysis of extracts from Rat heart, using TNFRSF14 Ab. The lane on the left was treated with blocking peptide.



DF6724 at 1/100 staining Human gastric cancer and adjacent normal tissues by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.