


IL10 Ab

[References\(45\)](#) [Images\(28\)](#)

Cat.#: DF6894	Concn.: ~1mg/ml	Mol.Wt.: 19kDa
Size:	Source: Rabbit	Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human,Mouse,Rat	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human IL10, corresponding to a region within the internal amino acids.	
Uniprot:	P22301	
Description:	IL-10 is an anti-inflammatory cytokine that is produced by T cells, NK cells and macrophages (1,2). IL-10 initiates signal transduction by binding to a cell surface receptor complex consisting of IL-10 RI and IL-10 RII . Binding of IL-10 leads to the activation of Jak1 and Tyk2, which phosphorylates Stat-3 (1,3). The anti-inflammatory activity of IL-10 is due to its ability to block signaling through other cytokine receptors, notably IFN γ receptor, by upregulating expression of SOCS-1 (1,3). In addition, IL-10 promotes T cell tolerance by inhibiting tyrosine phosphorylation of CD28 (4,5). IL-10 is an important negative regulator of the immune response, which allows for maintenance of pregnancy . In contrast, increased IL-10 levels contribute to persistent Leishmania major infections .	



DF6894 at 1/100 staining Rat ovary tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

procedures. Not for resale without express authorization.