

BLK Ab

[Images\(1\)](#)

Cat.#: BF0124	Concn.: ~1mg/ml	Mol.Wt.: 58kDa
Size:	Source: Mouse	Clonality: Monoclonal
Application:	ELISA 1:10000, WB 1:500-1:2000, IF/ICC 1:200-1:1000, FCM 1:200-1:400 *The optimal dilutions should be determined by the end user.	
Reactivity:	Human	
Storage:	Mouse IgG1 in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.	
Purification:	Affinity-chromatography.	
Immunogen:	Purified recombinant fragment of human BLK expressed in E. Coli.	
Uniprot:	P51451	
Description:	Blk is a Src family protein tyrosine kinase expressed in all stages of B cell development . Activation of B cells by various ligands is accompanied by activation of Blk . It has been suggested that Blk is involved in the control of B cell differentiation and proliferation . Blk transcripts have also been detected in human thymocytes, but not in mature T cells, implicating that Blk may play an important role in thymopoiesis . Blk function may be redundant, however, as mice that do not express Blk are not impaired with respect to B cell development and immune response .	




Figure 1: Western blot analysis using BLK mAb against HEK293 (1) and BLK(AA: 2-200)-hlgGfC transfected HEK293 (2) cell lysate.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.