

Affinity Biosciences website:www.affbiotech.com

RIPK3 Ab

References(1) Images(4)

Cat.#: DF7339 Mol.Wt.: 56kDa Concn.: ~1mg/ml Source: Rabbit Size: Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% Storage:

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

A synthesized peptide derived from mouse RIPK3, corresponding to a Immunogen:

region within the internal amino acids.

E9PZP3 Uniprot:

Description: The product of this gene is a member of the receptor-interacting protein

(RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is


predominantly localized to the cytoplasm, and can undergo

nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-

kappaB transcription factor.



Western blot analysis of extracts from B16F10, using RIPK3 Ab. The lane on the left was treated with blocking peptide.



DF7339 staining A549 cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF7339) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).



<code>TMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.