

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

CA9 Ab

Images(1)

| Cat.#: BF0375<br>Size:      | Concn.: ~1mg/ml<br>Source: Mouse  | Mol.Wt.: 50kDa<br>Clonality: Monoclonal |
|-----------------------------|---|---|
| Application:<br>Reactivity: | ELISA 1:10000, WB 1:500-1:2000, IHC 1:200-1:1000, FCM 1:200-1:400<br>*The optimal dilutions should be determined by the end user.<br>Human  |   |
| Storage:                    | Mouse IgG1 in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.   |   |
| Purification:               | Affinity-chromatography.  |   |
| Immunogen:                  | Purified recombinant fragment of human CA9 expressed in E. Coli.  |   |
| Uniprot:                    | Q16790  |   |
| Description:                | CA IX is a transmembrane protein and the only tumor-associated carbonic<br>anhydrase isoenzyme known. It is expressed in all clear-cell renal cell<br>carcinoma, but is not detected in normal kidney or most other normal<br>tissues. It may be involved in cell proliferation and transformation.<br>Reversible hydration of carbon dioxide. Participates in pH regulation. May<br>be involved in the control of cell proliferation and transformation. Appears<br>to be a novel specific biomarker for a cervical neoplasia. Tissue specificity:<br>Expressed primarily in carcinoma cells lines. Expression is restricted to very<br>few normal tissues and the most abundant expression is found in the<br>epithelial cells of gastric mucosa. |   |




Figure 1: Western blot analysis using CA9 mouse mAb against Hela (1) and A549 (2) cell lysate.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.