## Affinity Biosciences website:www.affbiotech.com

## **GAPDH Ab**

References(44) Images(28)

Cat.#: AF0911 Concn.: ~1mg/ml Mol.Wt.: 37KD
Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:5000-1:50000, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat, Pig, Zebrafish, Monkey

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink<sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).


Immunogen: A synthesized peptide derived from human GAPDH.

Uniprot: P04406

Description: Glyceraldehyde 3 phosphate dehydrogenase (GAPDH) is well known as one

of the key enzymes involved in glycolysis. GAPDH is constitutively abundant expressed in almost cell types at high levels, therefore antibodies against GAPDH are useful as loading controls for Western Blotting. Some pathology factors, such as hypoxia and diabetes, increased or decreased

GAPDH expression in certain cell types.




Western blot analysis of extracts from various samples, using GAPDH Ab.

Lane 1: Rat heart, blocked with antigen-specific peptides,

Lane 2: Rat heart, Lane 3: Mouse kidney,

Lane 4: A549 cells,

Lane 5: Hela cells(H2O2 treatment).



AF0911 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF0911) and mouse anti-beta tubulin Ab(T0023) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI (blue).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at  $4^{\circ}$ C with gentle shaking,



overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.