

## Affinity Biosciences website:www.affbiotech.com

## **Factor VIII Ab**

Images(2)

Cat.#: AF0156 Concn.: ~1mg/ml Mol.Wt.: 300kDa Size: Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:3000, IHC 1:50-1:200, IF/ICC 1:100-1:500

\*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLink<sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human Factor VIII, corresponding to a

region within C-terminal amino acids.

Uniprot: P00451


Description: F8 Factor VIII, along with calcium and phospholipid, acts as a cofactor for

factor IXa when it converts factor X to the activated form, factor Xa. Belongs to the multicopper oxidase family. Interacts with vWF. vWF

binding is essential for the stabilization of F8 in circulation.



Western blot analysis of extracts from Hela, using Factor VIII Ab.



AF0156 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.



For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.